

Research papers using Viewbox for cephalometric analysis (partial list):

1. Baccetti T, Antonini A, Franchi L, Tonti M, Tollaro I. Glenoid fossa position in different facial types: a cephalometric study. *Br J Orthod* 1997;24:55-9.
2. Baccetti T, Franchi L. Shape-coordinate and tensor analysis of skeletal changes in children with treated Class III malocclusions. *Am J Orthod Dentofacial Orthop* 1997;112:622-633.
3. Franchi L, Baccetti T, McNamara JA. Shape-coordinate analysis of skeletal changes induced by rapid maxillary expansion and facial mask therapy. *Am J Orthod Dentofacial Orthop* 1998;114:418-426.
4. Kazandjian S, Sameshima GT, Champlin T, Sinclair PM. Accuracy of video imaging for predicting the soft tissue profile after mandibular set-back surgery. *Am J Orthod Dentofacial Orthop* 1999;115:382-9.
5. Baccetti T, Franchi L, McNamara JA Jr. Thin-plate spline analysis of treatment effects of rapid maxillary expansion and face mask therapy in early Class III malocclusions. *Eur J Orthod* 1999;21:275-81.
6. Raadsheer MC, van Eijden TM, van Ginkel FC, Prah-Andersen B. Contribution of jaw muscle size and craniofacial morphology to human bite force magnitude. *J Dent Res* 1999;78:31-42.
7. Franchi L, Baccetti T, McNamara JA Jr. Thin-plate spline analysis of mandibular growth. *Angle Orthod* 2001;71:83-9.
8. Papadopoulos MA, Athanasiou AE, Peitsinis V, Ioannidou-Marathiotou I. Cephalometric evaluation of structural characteristics of mandibular rotation in individuals with Class III malocclusion and mandibular prognathism. *Hellenic Orthod Rev* 2001;4:81-102.
9. Baccetti T, Franchi L, McNamara JA Jr. An improved version of the cervical vertebral maturation (CVM) method for the assessment of mandibular growth. *Angle Orthod* 2002;72:316-23.
10. Faltin KJ, Faltin RM, Baccetti T, Franchi L, Ghiozzi B, McNamara JA Jr. Long-term effectiveness and treatment timing for Bionator therapy. *Angle Orthod* 2003;73:221-30.
11. Katsavrias EG. A method for integrating facial cephalometry and corrected lateral tomography of the temporomandibular joint. *Dentomaxillofac Radiol* 2003;32:93-6.
12. Katsavrias EG. The Effect of Mandibular Protrusive (Activator) Appliances on Articular Eminence Morphology. *Angle Orthod* 2003;73:647-53.
13. Halazonetis DJ. Morphometrics for cephalometric diagnosis. *Am J Orthod Dentofacial Orthop* 2004;125:571-81.

dHAL Software

14. Papadopoulos MA, Mavropoulos A, Karamouzos A. Cephalometric changes following simultaneous first and second maxillary molar distalization using a non-compliance intraoral appliance. *J Orofac Orthop* 2004;65:123-36.
15. Mavropoulos A, Bresin A, Kiliaridis S. Morphometric analysis of the mandible in growing rats with different masticatory functional demands: adaptation to an upper posterior bite block. *Eur J Oral Sci* 2004;112:259-266.
16. Katsavrias EG, Halazonetis DJ. Condyle and fossa shape in Class II and Class III skeletal patterns: a morphometric tomographic study. *Am J Orthod Dentofacial Orthop* 2005;128:337-46.
17. Mavropoulos A, Karamouzos A, Kiliaridis S, Papadopoulos MA. Efficiency of noncompliance simultaneous first and second upper molar distalization: a three-dimensional tooth movement analysis. *Angle Orthod.* 2005;75:532-9.
18. Leonardi M, Armi P, Baccetti T, Franchi L, Caltabiano M. Mandibular growth in subjects with infraoccluded deciduous molars: a superimposition study. *Angle Orthod* 2005;75:927-34.
19. Antonini A, Marinelli A, Baroni G, Franchi L, Defraia E. Class II malocclusion with maxillary protrusion from the deciduous through the mixed dentition: a longitudinal study. *Angle Orthod* 2005;75:980-6.
20. Ioannidou-Marathiotou I, Papadopoulos MA. Assessment of cephalometric analyses used for craniofacial complex evaluation: Part I - Sagittal maxillary position. *Hellenic Orthod Rev* 2005;8:15-32.
21. Camporesi M, Franchi L, Baccetti T, Antonini A. Thin-plate spline analysis of arch form in a Southern European population with an ideal natural occlusion. *Eur J Orthod* 2006;28:135-40.
22. Franchi L, Baccetti T. Prediction of individual mandibular changes induced by functional jaw orthopedics followed by fixed appliances in Class II patients. *Angle Orthod* 2006;76:950-4.
23. Katsavrias EG. Morphology of the temporomandibular joint in subjects with Class II Division 2 malocclusions. *Am J Orthod Dentofacial Orthop* 2006;129:470-8.
24. Nasiopoulos AT, Athanasiou AE, Papadopoulos MA, Kolokithas G, Ioannidou I. Premolar root changes following treatment with the banded Herbst appliance. *J Orofac Orthop* 2006;67:261-71.
25. Singleton DA, Buschang PH, Behrents RG, Hinton RJ. Craniofacial growth in growth hormone-deficient rats after growth hormone supplementation. *Am J Orthod Dentofacial Orthop* 2006;130:69-82.
26. Halazonetis DJ. Morphometric evaluation of soft-tissue profile shape. *Am J Orthod Dentofacial Orthop* 2007;131:481-9.

27. Halazonetis DJ. Morphometric correlation between facial soft-tissue profile shape and skeletal pattern in children and adolescents. *Am J Orthod Dentofacial Orthop* 2007;132:450-7.
28. Andredaki M, Koumantanou A, Dorotheou D, Halazonetis DJ. A cephalometric morphometric study of the sella turcica. *Eur J Orthod* 2007;29:449-456.
29. Dicker G, Van Spronsen P, Van Schijndel R, van Ginkel F, Manoliu R, Boom H, Tuinzing DB. Adaptation of jaw closing muscles after surgical mandibular advancement procedures in different vertical craniofacial types: a magnetic resonance imaging study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007;103:475-82.
30. Reyneke JP, Bryant RS, Suuronen R, Becker PJ. Postoperative skeletal stability following clockwise and counter-clockwise rotation of the maxillomandibular complex compared to conventional orthognathic treatment. *Br J Oral Maxillofac Surg* 2007;45:56-64.
31. Stalpers MJ, Booij JW, Bronkhorst EM, Kuijpers-Jagtman AM, Katsaros C. Extraction of maxillary first permanent molars in patients with Class II Division 1 malocclusion. *Am J Orthod Dentofacial Orthop* 2007;132:316-23.
32. Oosterkamp BC, Remmelink HJ, Pruim GJ, Hoekema A, Dijkstra PU. Craniofacial, craniocervical, and pharyngeal morphology in bilateral cleft lip and palate and obstructive sleep apnea patients. *Cleft Palate Craniofac J* 2007;44:1-7.
33. Carrillo R, Rossouw PE, Franco PF, Opperman LA, Buschang PH. Intrusion of multiradicular teeth and related root resorption with mini-screw implant anchorage: a radiographic evaluation. *Am J Orthod Dentofacial Orthop* 2007;132:647-55.
34. Carrillo R, Buschang PH, Opperman LA, Franco PF, Rossouw PE. Segmental intrusion with mini-screw implant anchorage: a radiographic evaluation. *Am J Orthod Dentofacial Orthop* 2007;132:576.e1-6.
35. Abed GS, Buschang PH, Taylor R, Hinton RJ. Maturation and functional related differences in rat craniofacial growth. *Arch Oral Biol* 2007;52:1018-25.
36. Defraia E, Camporesi M, Marinelli A, Tollaro I. Morphometric investigation in the skulls of young adults. *Angle Orthod* 2008;78:641-6.
37. Provatidis CG, Georgiopoulos B, Kotinas A, McDonald JP. Evaluation of craniofacial effects during rapid maxillary expansion through combined *in vivo/in vitro* and finite element studies. *Eur J Orthod* 2008;30:437-48.
38. Dvortsin DP, Sandham A, Pruim GJ, Dijkstra PU. A comparison of the reproducibility of manual tracing and on-screen digitization for cephalometric profile variables. *Eur J Orthod* 2008;30:586-91.
39. Bills GC, Buschang PH, Ceen R, Hinton RJ. Timing effects of growth hormone supplementation on rat craniofacial growth. *Eur J Orthod* 2008;30:153-62.

40. Chatzigianni A, Halazonetis DJ. Geometric morphometric evaluation of cervical vertebrae shape and its relation to skeletal maturation. *Am J Orthod Dentofacial Orthop* 2009;136:481.e1-481.e9.
41. Baccetti T, Franchi L, Stahl F. Comparison of 2 comprehensive Class II treatment protocols including the bonded Herbst and headgear appliances: a double-blind study of consecutively treated patients at puberty. *Am J Orthod Dentofacial Orthop* 2009;135:698.e1-10.
42. Staudt CB, Kiliaridis S. A nonradiographic approach to detect Class III skeletal discrepancies. *Am J Orthod Dentofacial Orthop* 2009;136:52-8.
43. Mayer Y, Machtei EE. Divergence correction associated with implant placement: a radiographic study. *Int J Oral Maxillofac Implants* 2009;24:1033-9.
44. Abed Y, Har-Zion G, Redlich M. Lip posture following debonding of labial appliances based on conventional profile photographs. *Angle Orthod* 2009;79:235-9.
45. van Vlijmen OJ, Bergé SJ, Swennen GR, Bronkhorst EM, Katsaros C, Kuijpers-Jagtman AM. Comparison of cephalometric radiographs obtained from cone-beam computed tomography scans and conventional radiographs. *J Oral Maxillofac Surg* 2009;67:92-7. doi: 10.1016/j.joms.2008.04.025.
46. van Vlijmen OJ, Bergé SJ, Bronkhorst EM, Swennen GR, Katsaros C, Kuijpers-Jagtman AM. A comparison of frontal radiographs obtained from cone beam CT scans and conventional frontal radiographs of human skulls. *Int J Oral Maxillofac Surg* 2009;38:773-8. doi: 10.1016/j.ijom.2009.02.024.
47. van Vlijmen OJ, Maal TJ, Bergé SJ, Bronkhorst EM, Katsaros C, Kuijpers-Jagtman AM. A comparison between two-dimensional and three-dimensional cephalometry on frontal radiographs and on cone beam computed tomography scans of human skulls. *Eur J Oral Sci* 2009;117:300-5. doi: 10.1111/j.1600-0722.2009.00633.x.
48. Koletsis DD, Halazonetis DJ. Cervical vertebrae anomalies in orthodontic patients. A growth-based superimpositional approach. *Eur J Orthod* 2010;32:36-42.
49. van Vlijmen OJ, Maal T, Bergé SJ, Bronkhorst EM, Katsaros C, Kuijpers-Jagtman AM. A comparison between 2D and 3D cephalometry on CBCT scans of human skulls. *Int J Oral Maxillofac Surg* 2010;39:156-60. doi: 10.1016/j.ijom.2009.11.017.
50. De Clerck H, Cevidanes L, Baccetti T. Dentofacial effects of bone-anchored maxillary protraction: a controlled study of consecutively treated Class III patients. *Am J Orthod Dentofacial Orthop* 2010;138:577-81.
51. Liu SS, Kyung HM, Buschang PH. Continuous forces are more effective than intermittent forces in expanding sutures. *Eur J Orthod* 2010;32:371-80.

dHAL Software

52. Tsorovas G, Karsten AL. A comparison of hand-tracing and cephalometric analysis computer programs with and without advanced features--accuracy and time demands. *Eur J Orthod* 2010;32:721-8.
53. Damstra J, Huddleston Slater JJ, Fourie Z, Ren Y. Reliability and the smallest detectable differences of lateral cephalometric measurements. *Am J Orthod Dentofacial Orthop* 2010;138:546.e1-8.
54. Halazonetis DJ. Permutation method for evaluating topographical craniofacial correlations. *Am J Orthod Dentofacial Orthop* 2011;139:e211-e217.
55. Gkantidis N, Halazonetis DJ. Morphological integration between the cranial base and the face in children and adults. *J Anat* 2011;218:426-38.
56. Bartzela TN, Katsaros C, Bronkhorst EM, Rizell S, Halazonetis D, Kuijpers-Jagtman AM. A two-centre study on facial morphology in patients with complete bilateral cleft lip and palate at nine years of age. *Int J Oral Maxillofac Surg* 2011;40:782-789.
57. Gkantidis N, Halazonetis DJ, Alexandropoulos E, Haralabakis NB. Contrasting treatment strategies in hyperdivergent Class II div. 1. Is vertical dimension affected? *Am J Orthod Dentofacial Orthop* 2011;140:346-55.
58. Livas C, Halazonetis DJ, Booij JW, Katsaros, C. Extraction of maxillary first molars improves second and third molar inclination in Class II Division 1. *Am J Orthod Dentofacial Orthop* 2011;140:377-82.
59. Bastir M, Rosas A, Gunz P, Peña-Melian A, Manzi G, Harvati K, Kruszynski R, Stringer C, Hublin JJ. Evolution of the base of the brain in highly encephalized human species. *Nat Commun* 2011;2:588.
60. Shaheed S, Iftikhar A, Rasool G, Bashir U. Accuracy of linear cephalometric measurements with scanned lateral cephalograms. *Pakistan Oral & Dental Journal* 2011;31:68-72.
61. Bartzela T, Katsaros C, Rønning E, Rizell S, Semb G, Bronkhorst E, Halazonetis D, Kuijpers-Jagtman AM. A longitudinal three-center study of craniofacial morphology at 6 and 12 years of age in patients with complete bilateral cleft lip and palate. *Clin Oral Investig* 2012;16:1313-24.
62. Thareja V, Shivaprakash G, Shamnur N, Kumar GA. An appraisal of indian profile attractiveness using digital image morphing. *J Ind Orthod Soc* 2012;46:9-16.
63. Johner AM, Pandis N, Dudic A, Kiliaridis S. Quantitative comparison of 3 enamel-stripping devices in vitro: how precisely can we strip teeth? *Am J Orthod Dentofacial Orthop* 2013;143(4 Suppl):S168-72. doi: 10.1016/j.ajodo.2012.10.001.
64. Livas C, Halazonetis DJ, Booij JW, Pandis N, Tu YK, Katsaros, C. Maxillary sinus floor extension and posterior tooth inclination in adolescent patients with

- Class II Division 1 malocclusion treated with maxillary first molar extractions
Am J Orthod Dentofacial Orthop 2013;143:479-85.
65. Wellens HLL, Kuijpers-Jagtman AM, Halazonetis DJ. Geometric morphometric analysis of craniofacial variation, ontogeny and modularity in a cross-sectional sample of modern humans. *J Anat* 2013;222:397-409.
 66. Polychronis G, Christou P, Mavragani M, Halazonetis DJ. Geometric morphometric 3D shape analysis and covariation of human mandibular and maxillary first molars. *Am J Phys Anthropol* 2013;152(2):186-96.
 67. Coquerelle M, Prados-Frutos JC, Rojo R, Mitteroecker P, Bastir M. Short faces, big tongues: developmental origin of the human chin. *PLoS One* 2013;8:e81287. doi: 10.1371/journal.pone.0081287.
 68. Valla K, Halazonetis DJ. Correlation of 2D:4D digit ratio and craniofacial shape in prepubertal children. *Am J Hum Biol* 2014;26(3):337-46.
 69. Polychronis G, Halazonetis DJ. Shape covariation between the craniofacial complex and first molars in humans. *J Anat* 2014;225:220-31.
 70. Franchi L, Pavoni C, Cerroni S, Cozza P. Thin-plate spline analysis of mandibular morphological changes induced by early class III treatment: a long-term evaluation. *Eur J Orthod* 2014;36:425-30.
 71. Danz JC, Greuter C, Sifakakis I, Fayed M, Pandis N, Katsaros C. Stability and relapse after orthodontic treatment of deep bite cases-a long-term follow-up study. *Eur J Orthod* 2014;36:522-30. doi: 10.1093/ejo/cjs079.
 72. Perinetti G, Caprioglio A, Contardo L. Visual assessment of the cervical vertebral maturation stages: A study of diagnostic accuracy and repeatability. *Angle Orthod* 2014;84:951-6. doi: 10.2319/120913-906.1.
 73. Bastir M, García-Martínez D, Estalrich A, García-Tabernero A, Huguet R, Ríos L, Barash A, Recheis W, de la Rasilla M, Rosas A. The relevance of the first ribs of the El Sidrón site (Asturias, Spain) for the understanding of the Neandertal thorax. *J Hum Evol* 2015; pii: S0047-2484(14)00248-6. doi: 10.1016/j.jhevol.2014.10.008.
 74. Perinetti G, Sbardella V, Contardo L. Diagnostic reliability of the third finger middle phalanx maturation (MPM) method in the identification of the mandibular growth peak. *Eur J Orthod* 2016. doi: <http://dx.doi.org/10.1093/ejo/cjw059>.
 75. Karell MA, Langstaff H, Halazonetis DJ, Minghetti C, Frelat M, Kranioti EF. A novel method for pair-matching using three-dimensional digital models of bone: mesh-to-mesh value comparison. *Int J Legal Med* 2016;130:1315-22. doi: 10.1007/s00414-016-1334-3.
 76. Cocos A, Halazonetis DJ. Craniofacial shape differs in patients with tooth agenesis: geometric morphometric analysis. *Eur J Orthod* 2016 doi: 10.1093/ejo/cjw049

77. Parcha E, Bitsanis E, Halazonetis DJ. Morphometric covariation between palatal shape and skeletal pattern in children and adolescents: a cross-sectional study. *Eur J Orthod* 2016 doi: 10.1093/ejo/cjw063
78. Katsadouris A, Halazonetis DJ. Geometric morphometric analysis of craniofacial growth between the ages of 12 and 14 in normal humans. *Eur J Orthod* 2016
79. Perillo L, Vitale M, Masucci C, D'Apuzzo F, Cozza P, Franchi L. Comparisons of two protocols for the early treatment of Class III dentoskeletal disharmony. *Eur J Orthod* 2016;38:51-56.
80. Perinetti G, Rosso L, Riatti R, Contardo L. Sagittal and vertical craniofacial growth pattern and timing of circumpubertal skeletal maturation: a multiple regression study. *Biomed Res Int* 2016;2016:1728712.